

## **Dominique Glauser:** full publication list in February 2017

Hostettler L, Grundy L, Käser-Pébernard S, Wicky C, William Schafer R & **Glauser DA** (2017). The bright fluorescent protein mNeonGreen facilitates protein expression analysis *in vivo*. **G3** (Genes, Genetics, Genomes) (online doi: 10.1534/g3.116.G3 (Genes, Genetics, Genomes) (online doi: 10.1534/g3.116.037747).

**Glauser DA** & Goodman MB (2016). Molecules empowering animals to sense and respond to temperature in changing environments. **Current Opinion in Neurobiology** 41: 92-98 (online DOI:10.1016/j.conb.2016.09.006).

Schild LC & **Glauser DA** (2015).

Dual color neural activation and behavior control with chrimson and CoChR in *Caenorhabditis elegans*. **Genetics** 200: 1029-1034 (online doi: 10.1534/genetics.115.17795).

Schild LC, Zbinden L, Bell HW, Yu YV, Sengupta P, Goodman MB & **Glauser DA** (2014). The balance between cytoplasmic and nuclear CaM Kinase-1 signaling controls the operating range of noxious heat avoidance. **Neuron** 84: 983-996.

Yu YV, Bell HW, **Glauser DA**, Van Hooser SD, Goodman MB & Sengupta P (2014). CaMKI-Dependent regulation of sensory gene expression mediates experience-dependent plasticity in the operating range of a thermosensory neuron. **Neuron** 84: 919-926.

Bou Dib P, Gnägi B, Daly F, Sabado V, Tas D, **Glauser DA**, Peter Meister P & Nagoshi E (2014). A Conserved Role for p48 Homologs in Protecting Dopaminergic Neurons from Oxidative Stress. **PLoS Genetics** 10(10):e1004718 (DOI: 10.1371/journal.pgen.1004718).

**Glauser DA** (2014).

The multiplicity of alternative splicing decisions in *Caenorhabditis elegans* is linked to specific intronic regulatory motifs and minisatellites. **BMC Genomics** 15:354, doi: 10.1186/1471-2164-15-364.

**Glauser DA** (2013).

How and why *Caenorhabditis elegans* uses distinct escape and avoidance regimes to minimise exposure to noxious heat. **Worm** 2(4): doi:10.4161/worm.27285.

Rufener L, Bedoni N, Baur R, Rey S, **Glauser DA**, Bouvier J, Beech R, Sigel E & Puoti A (2013). acr-23 Encodes a monepantel-sensitive channel in *Caenorhabditis elegans*. **PLoS Pathogens** 9(8) e1003524. doi:10.1371/journal.ppat.1003524.

Schild LC & **Glauser DA** (2013).

Dynamic switching between escape and avoidance regimes reduces *Caenorhabditis elegans* exposure to noxious heat. **Nature Communications** 4: #2198, doi:10.1038/ncomms31

**Glauser DA**, Johnson BE, Aldrich RW & Goodman MB (2011). Intron alternative splicing coordination is essential for *Caenorhabditis elegans* slo-1 gene function. **Proc. Nat. Acad. Sci. (USA)** 108: 20790-20795.

Johnson BE, **Glauser DA**, Dan-Glauser ES, Halling DB, Aldrich RW & Goodman MB. (2011). Alternatively spliced doamins interact to regulate BK potassium channel gating. **Proc. Nat. Acad. Sci. (USA)** 108: 20784-20789.

Geffeney SL, Cueva JG, **Glauser DA**, Doll JC, Lee TH, Montoya M, Karanis S, Garakani AM, Pruitt BL & Goodman MB (2011). DEG/ENaC but not TRP channels are the major mechanoelectrical transduction channels in a *C. elegans* nociceptor. **Neuron** 71: 845-857.

**Glauser DA**, Chen WC, Agin R, MacInnis BL, Hellman AB, Garrity PA, Tan MW & Goodman MB (2011). Heat avoidance is regulated by transient receptor potential (TRP) channels and a neuropeptide signaling pathway in *Caenorhabditis elegans*. **Genetics** 188: 91-103.

Ryser S, **Glauser D**, Vigier M, Zhang YQ, Tachini P, Schlegel W, Durand P & Irminger-Finger I (2011). Gene expression of rat spermatogonia and Sertoli cells reveals signaling pathways from stem cells to niche and testicular cancer cells to surrounding stroma. **BMC Genomics** #29.

**Glauser DA** & Goodman MB (2010). Neuropeptides strike back. **Nature Neuroscience** 13: 528-529.

**Glauser DA** & Schlegel W (2009). The FoxO/Bcl-6/cyclin D2 pathway mediates metabolic and growth factor stimulation of proliferation in Min6 pancreatic β-cells. **J Recept Signal Transduct Res.** 29: 293-298.

Walters EM, Garcia-Serres R, Naik SG, Bourquin F, **Glauser DA**, Schürmann P, Huynh BH & Johnson MK (2009). Role of Histidine-86 in the Catalytic Mechanism of Ferredoxin: Thioredoxin Reductase. **Biochemistry** 48: 1016-1024.

**Glauser DA** & Schlegel W (2008). FoxO proteins in pancreatic β cells as potential therapeutic targets in diabetes (2008). **Expert Review of Endocrinology and Metabolism** 3: 175-185.

**Glauser DA** & Schlegel W (2007). The emerging role of FOXO transcription factors in pancreatic β cells. **J Endocrinol.** 193: 195-207.

Dai S, Friemann R, **Glauser DA**, Bourquin F, Manieri W, Schürmann P & Eklund H (2007). Structural Snapshots along the Reaction Pathway of Ferredoxin:Thioredoxin Reductase. **Nature** 448: 92-96.

**Glauser DA**, Brun T, Gauthier BR & Schlegel W (2007). Transcriptional response of pancreatic beta cells to metabolic stimulation:

large scale identification of immediate-early and secondary response genes. **BMC Mol Biol.** 8: #54.

**Glauser DA** & Schlegel W (2007).

Sequential actions of ERK1/2 on AP-1 transcription factor allow temporal integration of metabolic signals in pancreatic  $\beta$  cells. **FASEB J.** 21: 3240-3249.

**Glauser DA** & Schlegel W (2006).

Mechanisms of transcriptional regulation underlying temporal integration of signals. **Nucleic Acids Research** 34: 5175-5183.

Walters EM, Garcia-Serres R, Jameson GN, **Glauser DA**, Bourquin F, Manieri W, Schürmann P, Johnson MK & Huynh BH (2005). Spectroscopic characterization of site-specific [Fe(4)S(4)] cluster chemistry in ferredoxin:thioredoxin reductase: implications for the catalytic mechanism. **J Am Chem Soc** 127: 9612-9624.

**Glauser DA**, Bourquin F, Manieri W & Schurmann P (2004).

Characterization of ferredoxin:thioredoxin reductase modified by site-directed mutagenesis. **J Biol Chem** 279: 16662-16669.